



pISSN 2385-0442  
eISSN 2385-0922

## Natural History Sciences

<https://sisn.pagepress.org/index.php/nhs/index>

**Publisher's Disclaimer.** E-publishing ahead of print is increasingly important for the rapid dissemination of science. **Natural History Sciences** is, therefore, E-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear on a regular issue of the journal.  
E-publishing of this PDF file has been approved by the authors.

To cite this Article:

Battisti C, 2022 - **The western European hedgehog *Erinaceus europaeus* L. 1758 on San Pietro Island (southern Sardinia, Italy) and an updated review of its presence on the Italian small islands.** Natural History Sciences, Milano doi: 10.4081/nhs.2022.594 [Epub Ahead of Print]

**The western European hedgehog *Erinaceus europaeus* L. 1758 on San Pietro Island (southern Sardinia, Italy) and an updated review of its presence on the Italian small islands**

Corrado Battisti

‘Torre Flavia’ LTER (Long Term Ecological Research) Station, Protected Area Service, Città Metropolitana di Roma, Via G. Ribotta, 41, 00144 Rome, Italy.  
E-mail: c.battisti@cittametropolitanaroma.it

**Acknowledgments**

Luciano Durante, Mario and Gabriele Masnata, Monica Gallarati, Pierpaolo Battisti and Mrs. Granara provided indirect information about the presence of this species on the island. Giovanni Amori, Francesco M. Angelici, Massimo Brunelli, Spartaco Gippoliti, Armando Nappi and Julien Vittier provided further data and suggestions. Alessandro Zocchi reviewed the English style and language. Francesco Zullo drew up Figure 1. Marzio Zapparoli provided the draft of Figure 2. The Editor (Giorgio Chiozzi) and two anonymous reviewers provided further useful comments and suggestions.

**Abstract** – The current presence and local distribution of the western European hedgehog *Erinaceus europaeus* L. 1758, on San Pietro Island (Southern Sardinia, Italy) are here discussed for the period 2013-2021 with a short review of the species' occurrence on the small Italian islands. The species was found in the central-eastern sector of the island, which is characterised by a high urban and infrastructural density and an agricultural mosaic, both suitable ecological conditions favouring its presence. Most of the direct data were obtained from road kills. Despite the elusive behaviour of the species and the consequent possible underestimation, it can be assumed that the species shows a low density. In this respect, it might be interesting to investigate whether local limiting factors, either anthropogenic or natural might act on the hedgehog population. At a larger scale, the updated review of the small Italian islands shows the presence of the species on 13 islands, of which probably only Elba, Asinara and San Pietro (all  $>50 \text{ km}^2$ ) support established populations. Further research is needed to study anthropogenic origin, history of introduction, genetics, density, and medium-long-term viability of these small populations in each insular context.

**Key words:** eradication, introduction, road kills, small island, viability.

**Riassunto** – Il riccio europeo occidentale *Erinaceus europaeus* L. 1758, sull'isola di San Pietro (Sardegna meridionale, Italia): distribuzione locale e revisione aggiornata della sua presenza nelle piccole isole italiane.

Una indagine sulla presenza del Riccio europeo occidentale su un'isola circumsarda (San Pietro), condotta dal 2013 al 2021, ha evidenziato una distribuzione limitata al settore centro-orientale, caratterizzato sia da una più alta densità urbana e infrastrutturale, sia dalla presenza di mosaici agricoli, condizioni ecologiche sinantropiche che possono favorire la presenza di questa specie. La maggior parte dei dati originali è stata ottenuta da investimenti con autoveicoli. Benché la specie mostri un comportamento elusivo e i dati potrebbero essere sottostimati, si può ipotizzare una bassa densità di popolazione sull'isola. A tal proposito, potrebbe essere interessante verificare se ciò possa essere fatto risalire a fattori locali limitanti, sia antropici sia naturali. Su scala nazionale, la revisione della letteratura evidenzia la presenza su 13 piccole isole italiane, con presenza di popolazioni stabili probabilmente solo su Elba, Asinara e San Pietro (tutte con superficie  $> 50 \text{ km}^2$ ). Ulteriori ricerche sono necessarie per studiare l'origine, la storia dell'introduzione, la genetica, la densità e la vitalità a medio-lungo termine di queste popolazioni in ciascun contesto insulare.

**Parole chiave:** eradicazione, introduzione, piccole isole, road kill, vitalità.

## INTRODUCTION

The western European hedgehog *Erinaceus europaeus* L. 1758 (Mammalia: Erinaceomorpha) is a common generalist species, widely distributed in mainland Italy, in Sicily, in Sardinia and in some small islands (Amori, 1993; Angelici *et al.*, 2009; Reggiani & Filippucci, 2008; Loy *et al.*, 2019; Tab. 1). In Sardinia, this mammal has been introduced in the middle and late neolithic (4<sup>th</sup> Millennium B.C.; Cetti, 1774; Cei, 1941; Sanges & Alcover, 1980; Vigne 1992; Masseti, 1993; Wilkens, 2003; Baldino *et al.*, 2008). However, although the hedgehog has been studied in Sardinia (e.g., roadkills: De Montis *et al.*, 2017; parasites: Scala & Garippa, 1996; Poglayen *et al.* 2003; Fois *et al.*, 2006; see also Regione Sardegna, 2017), data on its presence and local distribution on the circum-Sardinian islands, where this insectivore has been introduced, are still lacking (Angelici *et al.*, 2009; Masseti, 2019).

In this paper, I reported some direct and indirect records of presence and data on the local distribution of the western European hedgehog on the Island of San Pietro, a circum-Sardinian medium-sized island (Sulcis archipelago).

The western European hedgehog has also been recorded on several other small islands along peninsular Italy and on neighbouring large islands (Sicily and Sardinia: e.g. Angelici *et al.*, 2009). In this regard, given the interest in the presence and introduction history of this erinaceomorph in these contexts, I carried out an up-to-date review of scientific and grey literature on its presence on Italian small islands, also providing preliminary considerations on its status (occurrences of only occasionally introduced individuals vs. hypothetical stable populations).

## STUDY AREA

### San Pietro island

San Pietro Island (39°08'N; 8°18'E) is situated 11 km off the South-Western coast of Sardinia, opposite the Sulcis Peninsula. With a surface area of 51 km<sup>2</sup>, San Pietro Island is the sixth largest island in Italy (Fig. 1). The island is of volcanic origin (Cioni & Funedda, 2015; Gioncada *et al.*, 2019). Because of its high eco-biogeographic interest and the concern for its conservation (e.g., Arrigoni & Bocchieri, 1996; Bocchieri, 2001), San Pietro Island is listed as a Special Area of Conservation (code ITB040027 - SIC Isola di San Pietro; 92/43/EU Directive; Regione Sardegna, 2005).

The island is characterized mainly by a Mediterranean scrub (maquis with *Erica arborea* L., *Pistacia lentiscus* L., *Genista ephedroides* DC., *Juniperus turbinata* Guss. and *Arbutus unedo* L.). Mediterranean scrub is alternated with Mediterranean open prairies and anthropogenic pinewood patches (*Pinus pinaster* Aiton, *P. pinea* L., *P. halepensis* Mill.; further details in Regione Sardegna, 2015), these last widely diffused in the western sector.

After a long period in which the island was uninhabited, starting from 1738 the island was colonized by a human population of Ligurian origin that came from Tunisia (Vona *et al.*, 1996). In recent decades, the island's landscape has been progressively anthropized, with an increase in rural urbanisation, mainly in the eastern sector. In fact, the population is mostly concentrated in Carloforte (about 6,000 inhabitants), the only town on the island (Pellerano & Rivano, 1997; Pellerano, 2001). Climate is typically Mediterranean. A number of zoological studies have been carried out in this island in the last decades (see Baccetti *et al.*, 1989; Baccetti, 1996; for homeothermic vertebrates: bats: Zava *et al.*, 1995; birds: e.g., Schenk & Torre, 1988; Baccetti *et al.*, 1991; Rosén *et al.*, 1999; Battisti, 2018; Battisti & Zullo, 2019; Battisti & Fanelli, 2021).

### Italian small islands

Along the coasts of the Italian peninsula and large islands (Sicily and Sardinia) there are many small islands, sometimes grouped in archipelagos, ranging from 224 km<sup>2</sup> (Elba) to less than 1 km<sup>2</sup>. For this review, I included most of the Italian small islands, considering “small island” those “with limited surface and lack of environmental diversity” (La Greca & Sacchi, 1957). Small islands show different geological origin (sedimentary, metamorphic, volcanic effusive, volcanic intrusive). Most of the islands surveyed are situated relatively close to the mainland (excluding Pelagie Islands, located >70 km offshore). Human population in small islands ranges from 0 to >50,000 inhabitants (Ischia; for a review of their characteristics see Pretto *et al.*, 2012).

## METHODS

### San Pietro Island

In the 2013-2021 period, during holiday trips carried out in the summer season (July or August; approximately 20 days/year), I walked many pedestrian paths and paved roads (speed: 1.5 km/hour), covering all the main habitat types of the island (2013-2016: on an occasional basis; 2017-2021: >50 km/year and about 20 days/years of direct sampling; total: approximately >150 days of sampling effort). Interviews with local inhabitants were also conducted in an informal fashion. The collected data were all geo-referenced (latitude and longitude by Google Earth software ±10 m) and divided into direct (original) observations (living individuals and road-kills) and indirect records (obtained by interviewing local inhabitants or by web platforms: naturamediterraneo.com; ornitho.it; therio.it; iNaturalist.org), although with a lower degree of reliability.

### Italian small islands

In order to carry out a brief review of the presence of the western European hedgehog on the Italian small islands, I considered only small continental islands (see Watson, 2009; Ali, 2018; i.e. islands that had a connection to the mainland and/or to the larger islands or located <100 km away, such as the circum-peninsular, circum-Sardinian and circum-Sicilian islands). I excluded from the study those islands located in inland water basins (e.g., the Polvese Island, Lake Trasimeno, central Italy, where this species is present: Barzigotti & Capuano, 2006; Gaggi & Paci, 2014). To carry out the review, I considered all (at the best of my knowledge) scientific papers, grey literature and web platforms (naturamediterraneo.com; ornitho.it; therio.it; iNaturalist.org), obtaining updated information about the status of the species and on the viability of its populations on the small islands in the short and medium term (Table 2).

## RESULTS AND DISCUSSION

### San Pietro Island

During the study period, I obtained 15 records (7 direct observations, and 6 indirect records of 8 individuals) from 12 sites (Fig. 1, Tab. 1), mainly located in the central-eastern sector of the island, characterized by a higher urban and infrastructural density and agricultural mosaics (see Battisti, 2018; Battisti & Zullo, 2019; Battisti *et al.*, 2021), suitable ecological conditions that can favour the presence of this species (Reggiani & Filippucci, 2008). The lack of observations in the western sector may also be linked to the widespread presence of monospecific *Pinus pinaster* forests, which typically have poor undergrowth and therefore represent habitats of low ecological suitability for this species. (Reggiani & Filippucci, 2008).

Most of the direct data (6 out of 7 records; 85.7%) were obtained from roadkills (this species is highly sensitive to this anthropogenic impact: Rondinini & Doncaster, 2002; Battisti *et al.*, 2012; Wright *et al.*, 2020; see, for Sardinia: Amori *et al.*, 2014; De Montis *et al.*, 2017). However, despite considerable research effort, I have recorded low numbers of both road kills (even with the relatively high volume of vehicles during the summer period), and indirect detections of live animals, when compared to continental contexts (e.g., Reggiani & Filippucci, 2008). Although the species shows an elusive behaviour (Bearman-Brown *et al.*, 2020) and the data could be affected by underestimation, it can be hypothesized a sporadic presence on the island, with low densities. In this regard, it could be interesting to verify the role of indirect anthropogenic factors (e.g. road kill: Locatelli & Paolucci, 1998; Amori *et al.*, 2014; feral dogs: Reggiani & Filippucci, 2008) or natural processes, controlling this local population as, for example, the occasional predation by migrant and sedentary raptors, such as the barn owl *Tyto alba* (Mazzetti & Bortolotti, 1999) and common buzzard *Buteo buteo* (Swann & Etheridge, 1995, Sidorovich *et al.*, 2016), all locally present (Meschini & Frugis, 1993; Battisti, 2018; M. Masnata, pers. comm.). However, this is only hypothetical, since these two birds occur on the island with few individuals and their predation on hedgehog is unlikely. Also, the scarcity of both the hedgehog's preferred food items (e.g. earthworms and land snails: Reggiani & Filippucci, 2008) and of reliable water sources in this insular ecosystem might represent further local limiting factors.

About the origin of this insular population, there are anecdotal data on a probable recent introduction (presumably in the second half of the 20<sup>th</sup> century: M. Masnata, pers. comm.) from the neighbouring Sardinian mainland (this species may be easily introduced also involuntarily: see Nappi *et al.*, 2020). In this regard, further research is necessary.

The hedgehog has no conservation problems (Gippoliti & Amori, 2004; Bertolino *et al.*, 2015; para-autochthonous species: Senato della Repubblica, 2016), and does not represent a factor of local impact on species of conservation concern, especially if considering its low density on San Pietro Island. However, in other environmental contexts (for example, in wetland and coastal sites) evidence is shown of an impact of this mammal preying on clutches of waders and seabirds (see Utley *et al.*, 1989; Monteiro *et al.*, 1996; Jackson and Green, 2000). San Pietro is a site of high importance for waders, seabirds and other migratory and nesting species (Important Bird Area IT191; BirdLife International, 2001): in this regard, if further data will evidence circumstances of predation, an eradication project (e.g. by translocation) could be foreseeable, as suggested by Baccetti & Gotti (2016) and Iannucci *et al.*, (2019) for Pianosa Island (see also Warwick *et al.*, 2006; Capizzi, 2020).

#### Italian small islands

At national scale, the updated bibliographic and web-based review suggests a presence of the species on at least 13 small Italian islands (Tab. 2), with an updated confirmed absence in Ponza (Latium; Capizzi *et al.*, 2012), Ischia and Capri (Nappi *et al.* 2007), Tremiti (Ministero dell'Ambiente, 2021), Egadi (Felten & Storch, 1970; Zava & Lo Valvo, 1990; Ozella *et al.*, 2016) and Pelagie Islands (Arpa Sicilia, 2008; see also Amori, 1993; Angelici *et al.*, 2009). Although we do not have data available about local density, probably this mammal occurs with stable populations only on Elba Island (historical records reported by Thiebaud De Berneaud, 1808), and perhaps on San Pietro and Asinara, all islands with an area >50 km<sup>2</sup>, which may allow its persistence in the short and medium term (see Sara, 1998). The occasional reports obtained for the other islands are probably to be referred to single or to very few specimens introduced by humans, also voluntarily (the species is locally kept as pet or consumed for food: see Cornalia, 1870; Reggiani & Filippucci, 2008), therefore with a low probability of long-term persistence. In this sense, an in-depth and updated analysis of the local status and of the recent human-driven introduction/colonization history, possible local extinctions, genetics, and peculiar behaviours (e.g. absence of hibernation as found in southern populations: Aloise *et al.*, 2003) for the individual demographic units in the small islands could be of interest, contributing to the disciplinary arena of anthropogenic historical biogeography (e.g. see Balletto, 1996; Sarà, 1998; Mazza *et al.*, 2013; Amori *et al.*, 2015).

## REFERENCES

- Agnelli P., 1998 – I mammiferi dell’isola di Pianosa. In: Museo Zoologico “La Specola” dell’Università degli Studi di Firenze (ed.). Studio tecnico-scientifico sull’Isola di Pianosa. *Rapporto interno per l’Ente Parco Nazionale dell’Arcipelago Toscano*: 48–50.
- Ali J.R., 2018. Islands as biological substrates: continental. *Journal of Biogeography*, 45: 1003–1018.
- Aloise G., Scaravelli D., Bertozzi M. & Cagnin M. 2003 – Abbondanza relativa del Riccio *Erinaceus europaeus* L. 1758 (Insectivora, Erinaceidae) in ambienti del sud e nord Italia. *Hystrix, Italian Journal of Mammalogy*, 14(Supplement): 109.
- Amori G., 1993 – Italian insectivores and rodents: extinctions and current status. *Supplementi di Ricerche di Biologia della Selvaggina*, 21: 115-134.
- Amori G. & Masseti M., 1996 – Does the occurrence of predators on Central Mediterranean islands affect the body size of micromammals? *Vie et Milieu*, 46: 205–211.
- Amori G., Luiselli L., Milana G. & Casula P., 2014 – Distribuzione, diversità e abbondanza di micromammiferi associati ad habitat forestali in Sardegna. Report tecnico. *CNR – Istituto per lo studio degli ecosistemi, Roma ed Ente Foreste della Sardegna, Cagliari*.
- Amori G., Pinna V.R., Sammuri G. & Luiselli L., 2015 – Diversity of small mammal communities of the Tuscan Archipelago: testing the effects of island size, distance from mainland and human density. *Journal of Vertebrate Biology*, 64(2): 161-166.
- Angelici F.M. & Laurenti A., 1994 – I Mammiferi delle piccole isole italiane: stato delle conoscenze. *I Congresso Italiano di Teriologia, Pisa*, Riassunti, p. 9.
- Angelici F.M., Laurenti A. & Nappi A., 2009 – A checklist of the mammals of small Italian islands. *Hystrix, Italian Journal of Mammalogy* (n.s.), 20: 3-27.
- Arpa Sicilia, 2008 – Atlante della Biodiversità della Sicilia: Vertebrati terrestri. Studi e Ricerche, 6. *Arpa Sicilia, Palermo*.
- Arrigoni P.V. & Bocchieri E., 1996 – Caratteri fitogeografici della flora delle piccole isole circumsarde. *Biogeographia* 18: 63–90.
- Baccetti B., 1996 – Per una storia dell’esplorazione biogeografica delle isole che circondano la Sardegna. *Lavori della Società Italiana di Biogeografia* (n.s.), 18: 1–25.
- Baccetti B., Cobolli-Sbordoni M. & Poggi R., 1989 – Ricerche zoologiche della nave oceanografica “Minerva” (C.N.R.) sulle isole circumsarde. I. Introduzione. *Amali Museo Civico di Storia naturale Genova*, 87: 127-136.
- Baccetti N., Gotti C., 2016 – Protocollo di cattura e traslocazione del Riccio (*Erinaceus europaeus*) dall’Isola di Pianosa. Progetto LIFE13 NAT/IT/000471 “RESTO CON LIFE”. [Available on: [https://www.restoconlife.eu/wordpress/wp-content/uploads/2015/06/PE\\_Riccia.pdf](https://www.restoconlife.eu/wordpress/wp-content/uploads/2015/06/PE_Riccia.pdf)] [Accessed September 2021]
- Baccetti N., Roselli A. & Sposimo P., 1991 – Efficienza di una breve indagine sull’avifauna nidificante in una piccola isola mediterranea. Atti V Convegno italiano di Ornitologia. *Supplementi di Ricerche di Biologia della Selvaggina*, 17: 53-57.
- Baldino B., Carenti G., Grassi E., Orgolesu T., Secchi F. & Wilkens B. 2008 – L’economia animale dal Medioevo all’Età moderna nella Sardegna nord-occidentale. *Sardinia, Corsica et Baleares Antiquae*, 6: 103- 155.
- Balletto E., 1996 – Biogeografia insulare oggi: le isole del Mediterraneo. *Biogeographia-The Journal of Integrative Biogeography*, 18(1): 147-183.
- Battisti C., 2018 – Synanthropic-dominated biomass in an insular landbird assemblage. *Community Ecology*, 19(3): 203-210.
- Battisti C. & Fanelli G., 2021 – Alien-dominated plant communities’ syntopic with seabird’s nests: evidence and possible implication from a Mediterranean insular ecosystem. *Ethology Ecology & Evolution*, 33: 543-552.

- Battisti C. & Zullo F., 2019 – A recent colonizer bird as indicator of human-induced change: Eurasian collared dove (*Streptopelia decaocto*) in a small Mediterranean island. *Regional Environmental Change*, 19: 2113-2121.
- Battisti C., Amori G., De Felici S., Luiselli L. & Zapparoli M. 2012 – Mammal road-killing from a Mediterranean area in central Italy: evidence from an atlas dataset. *Rendiconti Lincei*, 23(2): 217-223.
- Battisti C., Zullo F. & Fanelli G., 2021 – The road to invasion: fine-grained distribution and suitability model for *Carpobrotus* sp. pl., a plant invader on a small Mediterranean island. *Folia Geobotanica*, 56: 1-11.
- Bearman-Brown L.E., Wilson L.E., Evans L.C. & Baker P.J., 2020 – Comparing non-invasive surveying techniques for elusive, nocturnal mammals: a case study of the West European hedgehog (*Erinaceus europaeus*). *Journal of Vertebrate Biology*, 69(3): 20075.
- Bertolino S., Colangelo P., Mori E. & Capizzi D., 2015 – Good for management, not for conservation: an overview of research, conservation and management of Italian small mammals. *Hystrix, Italian Journal of Mammalogy* (n.s.), 26(1): 25–35.
- BirdLife International, 2001 – Important Bird Areas and potential Ramsar Sites in Europe. BirdLife International, Wageningen.
- Bocchieri E. 2001 – Endemismi e rarità tra la flora delle piccole isole della provincia di Cagliari (Sardegna). *Biogeographia* 22: 139–168.
- Burzigotti R. & Capuano M.C. (Eds.), 2006 – Zoom sull’ambiente – L’isola Polvese. *Provincia di Perugia, Servizio Protezione Ambientale e Parchi, Perugia*.
- Capizzi D., 2020 – A review of mammal eradication on Mediterranean islands. *Mammal Review*, 50(2): 124-135.
- Capizzi D., Mortelliti A., Amori G., Colangelo P. & Rondini C. (Eds.), 2012 – I mammiferi del Lazio. Distribuzione, ecologia e conservazione. *Edizioni ARP, Roma*.
- Cei, J.M.A.M., 1941 – Ipotesi e ricerche biogeografiche sulle grandi isole del Tirreno: Le conoscenze zoogeografiche odierne. *Rivista Geografica Italiana*, 48: 283-292.
- Cetti F., 1774 – I quadrupedi di Sardegna. *Sassari*.
- Cioni R. & Funedda A., 2005 – Structural geology of crystal-rich, silicic lava flows: a case study from San Pietro Island (Sardinia, Italy). *Special papers, Geological Society of America*, 396: 1.
- Cornalia E., 1870 – Fauna d’Italia. Parte prima. Catalogo descrittivo dei mammiferi osservati fino ad ora in Italia. *Vallardi*, Milano.
- Cossu A., Monballiu X. & Torre A., 1994 – L’isola dell’Asinara. *Delfino, Sassari*.
- Cristaldi M., Amori G., Cagnin M., De Angelis R., Degrassi F., De Sanctis P., Filippucci M.G., Ieradi L.A., Pacilli A.M., Paggi S., Salucci M.P., Tommasi M. & Van Axel Castelli I., 1987 – Interrelazioni tra popolamenti a mammiferi e fattori antropici negli ambienti insulari eoliani. In: Giavelli G. & Moroni A. (Eds.), L’approccio interdisciplinare nelle ricerche sull’arcipelago eoliano. *Istituto d’Ecologia Università di Parma*: 51-76.
- Damiani G., 1923 – La Fauna. L’Elba illustrata. *S. Foresi Editore*: 103-129.
- De Marinis A.M., Masseti M. & Sforzi A., 1996 – Note on the non-flying terrestrial mammals of the Tuscan Archipelago, Northern Tyrrhenian Sea (Italy). *Bollettino del Museo Regionale di Scienze Naturali, Torino*, 14 (1): 275-281.
- De Montis A., Ledda A., Serra V. & Peddio V., 2017 – Assessing landscape fragmentation in Sardinia (Italy): the barrier effect. In: *Geographic Information Systems Conference and Exhibition “GIS Odyssey 2017”*: 126.
- Ente Parco Nazionale dell’Arcipelago Toscano, 2016 – Piano del Parco. Allegato 1a. Agriconsulting S.p.A., Roma. available on: <http://www.parks.it/dbdoc/PDF/statuto539.pdf>
- Felten H. & Storch G., 1970 – Kleinsäuger von den italienischen Mittelmeer-Inseln Pantelleria und Lampedusa (Mammalia). *Senckenbergiana biologica*, 51: 159-173.

- Fois F., Piras P., Addis G., Cabras P., Patta C. & Satta G., 2006 – Identification of ticks collected in two areas of Sardinia. *Parassitologia* 48: 349.
- Gaggi A. & Paci A.M., 2014 – Atlante degli Erinaceomorfi, dei Soricomorfi e dei piccolo roditori dell’Umbria. *Regione Umbria, Perugia*.
- Gioncada A., Pitzalis E., Cioni R., Fulignati P., Lezzerini M., Mundula F. & Funedda A., 2019 – The volcanic and mining geoheritage of San Pietro Island (Sulcis, Sardinia, Italy): the potential for geosite valorization. *Geoheritage*, 11(4): 1567-1581.
- Gippoliti S. & Amori G., 2004 – Mediterranean Island mammals: are they a priority for biodiversity conservation? *Biogeographia–The Journal of Integrative Biogeography*, 25(1): 135-144.
- Iannucci A., Baccetti N., Giannini F., Gotti C. & Baratti M., 2019 – A genetic analysis of the European hedgehog (*Erinaceus europaeus*): an applicative case study to support its eradication from Pianosa Island (Tuscan Archipelago). *Conservation Genetics*, 20(2): 395-402.
- Jackson D.B. & Green R.E. 2000 – The importance of the introduced hedgehog (*Erinaceus europaeus*) as a predator of the eggs of waders (*Charadrii*) on machair in South Uist, Scotland. *Biological Conservation*, 93(3): 333-348.
- La Greca M. & Sacchi C.F., 1957 – Problemi del popolamento animale nelle piccole isole mediterranee. *Annuario dell’Istituto e Museo di Zoologia dell’Università di Napoli*, 9: 1-189.
- Lo Cascio P. & Navarra E., 2003 – Guida naturalistica alle isole Eolie. *L’Epos società editrice, Palermo*.
- Locatelli R. & Paolucci P., 1998 – Il Riccio (*Erinaceus europaeus* L., 1758) nel Parco regionale dei Colli Euganei: alimentazione e impatto del traffico stradale. *Atti 2° Convegno Faunisti Veneti, Supplementi del Bollettino del Museo Civico di Storia naturale di Venezia*, 48: 94-104.
- Loy A., Aloise G., Ancillotto L., Angelici F.M., Bertolino S., Capizzi D., Castiglia R., Colangelo P., Contoli L., Cozzi, B., Fontaneto D., Lapini L., Maio N., Monaco A., Mori E., Nappi A., Podestà M., Russo D., Sarà M., Scandura M. & Amori G., 2019 – Mammals of Italy: An annotated checklist. *Hystrix, Italian Journal of Mammalogy*, 30: 87–106.
- Masseti M., 1993 – Post-Pleistocene variations of the non-flying terrestrial mammals on some Italian islands. *Supplementi di Ricerche di Biologia della Selvaggina*, 21: 209-217.
- Masseti M., 2019 – Terrestrial mammals of the satellite islands of Sardinia (Italy). *Biodiversity Journal*, 10 (4): 373–382.
- Mazzotti S. & Bortolotti C., 1999 – Specie-preda come indicatori delle variazioni stagionali della dieta del Barbagianni, *Tyto alba*. *Rivista italiana di Ornitologia*, 69: 89-96.
- Mazza P.P., Lovari S., Masini F., Massetti M. & Rustioni M., 2013 – A multidisciplinary approach to the analysis of multifactorial land mammal colonization of islands. *BioScience*, 63(12): 939-951.
- Meschini E. & Frugis S. (Eds.), 1993 – Atlante degli uccelli nidificanti in Italia. *Supplementi di Ricerche di Biologia della Selvaggina*, 20.
- Ministero dell’Ambiente, 2021 – Area marina protetta ‘Isole Tremiti’. [Available on line: <https://www.naturaitalia.it/apriAreaNaturale.do?idAreaNaturale=36&numeroPuntoInformativo=3>] [Accessed 2 september 2021]
- Monteiro L.R., Ramos J.A. & Furness R.W., 1996 – Past and present status and conservation of the seabirds breeding in the Azores archipelago. *Biological Conservation*, 78(3): 319-328.
- Nappi A. & Massetti M., 2005 – I mammiferi terrestri non volatori delle isole del Golfo di Napoli. Atti V Congresso Italiano di Teriologia. *Hystrix, Italian Journal of Mammalogy* (n.s.), suppl. 2005: 44.
- Nappi A., Cipolla R.M., Gabriele R., Massetti M., Corti C. & Arcidiacono G., 2007 – Anfibi, Rettili e Mammiferi delle isole del Golfo di Napoli: check-list commentata. Atti XVII Convegno del Gruppo per l’Ecologia di Base “G. Gadio”. *Studi Trentini di Scienze Naturali, Acta Biologica*, 83: 93-97.

- Nappi A., Ferloni M., Bassi E., 2020 – Updated distribution of *Erinaceus europaeus* Linnaeus, 1758 (Mammalia, Erinaceomorpha) in the Province of Sondrio (Italy). *Rivista del Museo Civico di Scienze Naturali "Enrico Caffi", Bergamo*, 33: 15-22.
- Ozella L., Cecchetti M. & Pessani D., 2016 – Diet of feral cats during the Scopoli's shearwater breeding season on Linosa Island, Mediterranean Sea. *Italian Journal of Zoology*, 83(4), 589-599.
- Pellerano L. & Rivano A., 1997 – Natura dell'isola di San Pietro. La flora. *Edizioni della Torre, Cagliari*.
- Pellerano L., 2001 – L'ambiente marino e costiero dell'isola di San Pietro. *Edizioni della Torre, Cagliari*.
- Poglayen G., Giannetto S., Brianti E., Scala A., Garippa G., Capelli G., Scaravelli D. & Reeve N.J., 2003 – Helminthes found in hedgehogs (*Erinaceus europaeus*) in three areas of Italy. *Veterinary Record*, 152: 22–24.
- Presutti F., 2008 – Arcipelago di La Maddalena. La fauna. [Available on: <https://arcieplagolamaddalena.wordpress.com/2018/02/13/la-fauna/>] [Accessed September 2021]
- Pretto F., Celesti-Grapow L., Carli E., Brundu G. & Blasi C., 2012 – Determinants of non-native plant species richness and composition across small Mediterranean islands. *Biological Invasions*, 14(12): 2559-2572.
- Reggiani G. & Filippucci M.G., 2008 – *Erinaceus europaeus* Linnaeus, 1758. In: Amori G., Contoli L. & Nappi A. (Eds.), Fauna d'Italia, Mammalia II: Erinaceomorpha, Soricomorpha, Lagomorpha, Rodentia. *Edizioni Calderini, Bologna*: 70-81.
- Regione Sardegna, 2015 – Piano di gestione del SIC “ITB040027 Isola di San Pietro”. [Available online: <https://portal.sardegnasira.it/web/sardegnaambiente/rete-natura-2000> [Accessed February 2022]
- Regione Sardegna, 2017 – *Erinaceus europaeus italicus*. Sardegna Foreste. *Regione Autonoma della Sardegna*. <http://www.sardegnaambiente.it/j/v/159?c=1581&s=9046&t=1&v=2>.
- Rondinini C. & Doncaster C.P., 2002 – Roads as barriers to movement for hedgehogs. *Functional Ecology*, 16(4): 504-509.
- Rosén M., Hedenström A., Badami A., Spina F. & Åkesson S., 1999 – Hunting flight behaviour of the Eleonora's Falcon *Falco eleonorae*. *Journal of Avian Biology*, 30: 342-350.
- Ruffo S. & Stoch F. (eds.), 2005 – Checklist e distribuzione della fauna italiana. *Memorie del Museo Civico di Storia Naturale di Verona, 2<sup>a</sup> serie, Sezione Scienze della Vita*, 16.
- Sanges M. & Alcover J.A., 1980 – Noticia sobre la microfauna vertebrada holocenica de la Grotta Su Guanu o Gonagosula (Oliena, Sardenya). *Endins*, 7: 57-62.
- Sarà M., 1998 – I mammiferi delle isole del Mediteraneo. *L'Epos, Palermo*.
- Scala A. & Garippa G., 1996 – Principal helminthiases of the hedgehog (*Erinaceus europaeus*) in Sardinia: epidemiological anatomical and histopathological findings. *Atti del Convegno Nazionale: Ecopatologia della Fauna Selvatica, Bologna, Italy*, 15-17 dicembre 1994., 155-161.
- Scaravelli D., 1999 – The status of *Erinaceus europaeus italicus* and *Erinaceus (concolor) roumanicus* in Italy. In: Reeve N. (ed.), *Proceedings of III International Hedgehog Workshop of the European Hedgehog Research Group, London, 29/30 January 1999. Roehampton, London*: 19.
- Schenk H. & Torre A., 1988 – L'avifauna nidificante della fascia costiera della Sardegna, comprese le isole circumsarde, 1978-87. *Il Naturalista siciliano*, s. 4, 12: 187-192.
- Senato della Repubblica, 2016 – Dossier del Servizio Studi sull'A.S. n. 119-A "Nuove disposizioni in materia di aree protette", ottobre 2016, n. 396 [Available on: [https://www.senato.it/japp/bgt/showdoc/17/DOSSIER/0/993044/index.html?part=dossier\\_dossier1-sezione\\_sezione15](https://www.senato.it/japp/bgt/showdoc/17/DOSSIER/0/993044/index.html?part=dossier_dossier1-sezione_sezione15); Accessed September 2021].

- Spagnesi M., De Marinis A.M. (Eds.), 2002 – Mammiferi d’Italia. Quad. Cons. Natura, 14. Ministero dell’Ambiente, Istituto Nazionale per la Fauna Selvatica, Roma.
- Spagnesi M., Toso S. & De Marinis A.M. (Eds.), 2004 – Iconografia dei mammiferi d’Italia. Seconda edizione. *Istituto Nazionale per la Fauna selvatica, Ministero dell’ambiente e della Tutela del Territorio, Roma*.
- Sidorovich A.A., Ivanovskij V.V., Sidorovich V.E. & Solovej I.A., 2016 – Landscape-related variation in the diet composition of the common buzzard (*Buteo buteo*) in Belarus. *Slovak Raptor Journal*, 10(1): 65-74.
- Swann R.L. & Etheridge B., 1995 – A comparison of breeding success and prey of the Common Buzzard *Buteo buteo* in two areas of northern Scotland. *Bird Study*, 42(1): 37-43.
- Thibault J.C., Delaugerre M., Cheylan G., Guyot I. & Miniconi R., 1987 – Les vertébrés terrestres non domestiques des îles Lavezzi (sud de la Corse). *Bulletin de la Société Linnéenne de Lyon*, 56: 73– 103, 117–152.
- Thiebaud De Berneaud A., 1808 – Voyage à l’isle d’Elbe, suivi d’une notice sur les autres îles de la Mer Tyrrhenienne. *D. Colas & Le Normant, Paris*.
- Tomassone L., Grego E., Auricchio D., Iori A., Giannini F. & Rambozzi L., 2013 – Lyme borreliosis spirochetes and spotted fever group *rickettsiae* in ixodid ticks from Pianosa island, Tuscany Archipelago, Italy. *Vector-borne and Zoonotic Diseases*, 13(2), 84-91.
- Torre A. & Monbailliu X., 1993 – La Fauna. In: Cossu A., Gazale V., Monbailliu X. & Torre A. (Eds.), Isole minori del Mediterraneo. Asinara. Storia, natura, mare e tutela ambientale. *Edizioni Delfino, Sassari*.
- Uttley J., Monaghan P. & Blackwood J., 1989 – Hedgehog *Erinaceus europaeus* predation on Arctic tern *Sterna paradisaea* eggs: the impact on breeding success. *Seabird*, 12: 3-6.
- Vesmanis I.E. & Hutterer R., 1980 – Nachweise von *Erinaceus*, *Crocidura* und *Microtus*, für die Insel Elba, Italien. *Zeitschrift für Säugetierkunde*, 45: 251-253.
- Vigne J. D., 1992 – Zooarchaeology and the biogeographical history of the mammals of Corsica and Sardinia since the last ice age. *Mammal Review*, 22(2), 87-96.
- Warwick H., Morris, P. & Walker D., 2006 – Survival and weight changes of hedgehogs (*Erinaceus europaeus*) translocated from the Hebrides to Mainland Scotland. *Lutra*, 49(2), 89-102.
- Watson D.M. 2009 – Continental islands. In: Gillespie R.G. & Clague D.A. (Eds), Encyclopedia of islands. *University of California Press*, pp. 180–187.
- Wilkens B., 2003 – La fauna sarda durante l’Olocene: le conoscenze attuali. *Sardinia, Corsica et Baleares antiquae: International Journal of Archaeology*: 1: 181-197.
- Wright, P.G., Coomber, F.G., Bellamy, C.C., Perkins, S.E. & Mathews, F. 2020 – Predicting hedgehog mortality risks on British roads using habitat suitability modelling. *PeerJ*, 7, e8154.
- Zava B. & Lo Valvo F., 1990 – First record of *Suncus etruscus* and notes on the bats of Pantelleria island, Italy. *Mammalia*, 54: 661-663.
- Zava B., Fiore M., Fornasari L. & Violani C., 1995 – Note sui Chiroterri dell’Isola di San Pietro con cenni storici sulle ricerche chiroterologiche in Sardegna. *Biogeographia*, 18: 641–651.

N	Year	Site and geographical coordinates	Notes (nr. ind., type of record, observer)
1	2015	Macchione (39°07'58"N, 8°18'00"E)	1 ind, roadkill (C. Battisti, pers. obs.)
2	2019	Girin (39°06'51"N, 8°18'25"E)	1 ind., roadkill (C. Battisti, pers. obs.)
3	2019	Valacca (39°07'25"N, 8°17'29"E)	2 ind. (P. Battisti, G. Masnata, pers. obs.)
4	2019 (8 August)	Macchione (39°07'54"N, 8°18'00"E)	1 ind., roadkill (C. Battisti, pers. obs.)
5	2019 (13 August)	Gioia, Sebino (39°08'42"N, 8°16'26"E)	1 ind., roadkill (C. Battisti, pers. obs.)
6	2021	La Punta, Memmerosso (39°10'13"N, 8°18'02"E)	1 ind., roadkill (M. Gallarati, pers. obs.)
7	Post 2000	Giunco (39°07'32"N, 8°18'30"E)	Adult and juveniles (M. Masnata, pers. obs.)
8	Post 2010	Valacca (39°07'24"N, 8°17'28"E), provincial road	Ind. near a fountain (P. Battisti and G. Masnata, pers. obs.)
9	Post 2000	Bacciu (39°07'38"N, 8°17'40"E)	Recorded by a dog in a private garden (M. Masnata, pers. obs.)
10	Post 2000	Girin (39°06'54"N, 8°18'25"E)	M. Masnata, pers. obs.
11	Post 2010	Punta del Morto, Carloforte (39°09'01"N, 8°18'37"E)	2 ind. (M. Masnata, pers. obs.)
12	2017 (7 October)	Valacca (39°07'17"N, 8°17'19"E), provincial road	1 ind., roadkill (Julien Vittier, pers. obs. reported on Ornitho.it)

Tab. 1 – Author's direct observations (1-6) and indirect records (7-12) of western European Hedgehog on San Pietro Island (2013-2021 period). For the indirect records (7-12), the reported coordinates are approximate. / Osservazioni dirette dell'autore (1-6) e segnalazioni indirette (7-12) di riccio europeo occidentale sull'Isola di San Pietro (periodo 2013-2021). Per le segnalazioni indirette (7,12), le coordinate riportate sono approssimative.

Small island (and size area)	Ref.	Notes	Probable stable population (p), occasional presence of individuals without evidence of stable populations (i)
Elba (224 km <sup>2</sup> )	Thiebaud De Berneaud, 1808; Damiani, 1923; Vesmanis & Hutterer 1980; De Marinis <i>et al.</i> , 1996; Amori & Masseti, 1996 (see also Amori, 1993; Sarà, 1998; Scaravelli, 1999; Spagnesi & De Marinis, 2002; Spagnesi <i>et al.</i> , 2004; Ruffo & Stoch, 2005; Arpa Sicilia, 2008; Angelici <i>et al.</i> , 2009; Ente Parco Nazionale dell'Arcipelago Toscano, 2016).	Documented presence for the last 200 years (Thiebaud De Berneaud, 1808); can be considered as native (see Iannucci <i>et al.</i> , 2009); 1 ind. roadkilled in Capoliveri (1993) (De Marinis <i>et al.</i> , 1996).	p
Capraia (19.3 km <sup>2</sup> )	Angelici <i>et al.</i> , 2009, and references therein.	2 ind. observed (summer 1992; F.M. Angelici pers. obs.) Species not included in the National Park plan (Ente Parco Nazionale Arcipelago Toscano, 2016). Not reported in Sarà (1998).	i
Pianosa (10.2 km <sup>2</sup> )	Agnelli, 1998; Ruffo & Stoch, 2005; Iannucci <i>et al.</i> , 2019 (see also Reggiani & Filippucci, 2008; Tomassone <i>et al.</i> , 2013; Ente Parco Nazionale dell'Arcipelago Toscano, 2016).	Population of Pianosa has an extremely low genetic diversity and a profile very similar to that of Elba (originated from a pool of individuals, perhaps also only a single pregnant female, moved by human from Elba in recent times; absent before 1950s Baccetti & Gotti, 2016; Iannucci <i>et al.</i> , 2019). Not reported in Sarà (1998).	i (p?)
Alicudi (5.2 km <sup>2</sup> )	Cristaldi <i>et al.</i> , 1987; Lo Cascio & Navarra, 2003 (see also Amori, 1993; Sarà, 1998; Scaravelli, 1999; Spagnesi & De Marinis, 2002; Spagnesi <i>et al.</i> , 2004; Arpa Sicilia, 2008; Angelici <i>et al.</i> , 2009).	Introduced in the second half of the 20 <sup>th</sup> century (beginning of the 1980s; Cristaldi <i>et al.</i> , 1987), 1 ind. observed in loc. Filo dell'Arpa (photo available; Lo Cascio & Navarra, 2003; see also Arpa Sicilia, 2008).	i
Salina (26.1 km <sup>2</sup> )	Lo Cascio & Navarra, 2003.	1 ind. in a private garden in first 2000s (Lo Cascio & Navarra, 2003).	i
Favignana (19.8 km <sup>2</sup> )	Sarà, 1998 (see also Arpa Sicilia, 2008; Reggiani & Filippucci, 2008; Angelici <i>et al.</i> 2009).	Occasional presence with a specimen found dead (roadkill) in 1989 (Sarà, 1998). Recently introduced (Arpa Sicilia, 2008).	i
Ustica (8.7 km <sup>2</sup> )	Sarà, 1998 (see also Reggiani & Filippucci, 2008; Angelici <i>et al.</i> 2009).	1 ind. observed for a few months (1994) in a cropland (Sarà, 1998).	i
Procida (4.1 km <sup>2</sup> )	Nappi & Masseti, 2005 (and references therein); Nappi <i>et al.</i> , 2007 (see also Reggiani & Filippucci, 2008; Angelici <i>et al.</i> 2009;	Established in the last years (Nappi & Masseti, 2005).	i

	www.isoledellacampania.eu/procida-1.html).		
Vivara (0.4 km <sup>2</sup> ) <sup>(*)</sup>	M. Rodriguez (pers. obs.) cit. in Nappi <i>et al.</i> , 2007 (see also Angelici <i>et al.</i> 2009).	Reported for the second half of 90s. Data are not sufficient to confirm the existence of a stable population (Nappi <i>et al.</i> , 2007). Not reported in Sarà (1998).	i
Caprera (16 km <sup>2</sup> )	Torre & Maiabailiu, 1993; Cossu <i>et al.</i> , 1994 (see also Angelici <i>et al.</i> 2009; Presutti, 2018; Masseti, 2019).	Recently introduced in Caprera (Cossu <i>et al.</i> , 1994).	i
La Maddalena (20 km <sup>2</sup> )	Angelici <i>et al.</i> , 2009, and references therein.		i?
Asinara (52 km <sup>2</sup> )	Torre & Maiabailiu, 1993; Cossu <i>et al.</i> , 1994; Amori & Masseti, 1996 (see also Sarà, 1998; Spagnesi & De Marinis, 2002; Spagnesi <i>et al.</i> , 2004; Ruffo & Stoch, 2005; Arpa Sicilia, 2008; Reggiani & Filippucci, 2008; Angelici <i>et al.</i> 2009; Masseti, 2019).	Recently introduced (Cossu <i>et al.</i> , 1994).	p?
San Pietro (51 km <sup>2</sup> )	Zava <i>et al.</i> , 1995 (see also Ruffo & Stoch, 2005; Reggiani & Filippucci, 2008; Masseti, 2019). This study.	Recorded in 1989 (Zava <i>et al.</i> , 1995).	p?

Tab. 2 – The western European Hedgehog on the small Italian islands. Bibliographic and web-based evidence of presence for each small island, notes and considerations about probable occurrence of stable populations (p) *vs.* only occasional presence of individuals (i) have been reported. Area: size area (in km<sup>2</sup>). (\*): Vivara is connected to Procida by a bridge. / Il riccio europeo occidentale nelle piccole isole italiane. Per ciascuna piccola isola, sono stati riportati sia i riferimenti bibliografici (e sitografici) che segnalano la specie, sia note e considerazioni sulla probabile presenza di popolazioni stabili (p) piuttosto che la sola presenza occasionale di individui (i). Area: superficie dell'isola (in km<sup>2</sup>). (\*): Vivara è collegata a Procida da un ponte.




Fig. 1 – The study area (San Pietro Island, Sulcis archipelago; South Sardinia) with the evidence of presence of western European hedgehog. Black circles: direct observations; Grey circles: indirect data. See Table 1 for details and numbers. / L'area di studio (Isola di San Pietro, arcipelago del Sulcis; Sardegna meridionale) con i siti di presenza del riccio europeo occidentale. Cerchi neri: osservazione diretta; cerchi grigi: dati indiretti. Vedere Tabella 1 per dettagli e numerazione.




Fig. 2 – The small Italian islands with evidence of western European hedgehog presence. Thick-bordered circles: probable occurrence of stable populations; thin-bordered circles: only occasional presence of individuals (low probability of short-term persistence). See Table 2 for details. 1: Elba; 2: Capraia; 3: Pianosa; 4: Alicudi; 5: Salina; 6: Favignana; 7: Ustica; 8: Procida; 9: Vivara; 10: Capraia; 11: La Maddalena; 12: Asinara; 13: San Pietro. / Evidenze di presenza di riccio europeo occidentale nelle piccole isole italiane. Cerchi con bordo spesso: probabile presenza di popolazioni stabili; cerchi con bordo sottile: probabile presenza di singoli individui (bassa probabilità di persistenza a breve termine). Vedasi Tabella 2 per dettagli: 1: Elba; 2: Capraia; 3: Pianosa; 4: Alicudi; 5: Salina; 6: Favignana; 7: Ustica; 8: Procida; 9: Vivara; 10: Capraia; 11: La Maddalena; 12: Asinara; 13: San Pietro.